

Original Article



Can 5-minute Apgar score be considered as an additional independent risk factor for screening of retinopathy of prematurity?

Amir Eftekhari Milani¹, Mohamad Reza Niyousha¹, Ali Kiavar^{1*}, Hanieh Sakha², Ali Mahdavi Fard³, Zahra Abdollahi²

¹Department of Ophthalmology, Tabriz University of Medical Sciences, Tabriz, Iran.

²School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

³Department of Ophthalmology, University at Buffalo, Buffalo, NY, USA.

Article info

Article History:

Received: 1 Nov. 2020

Accepted: 26 Jan. 2020

e-Published: 7 March 2020

Keywords:

- Apgar score
- Prematurity
- Retinopathy

Abstract

Introduction: Because of increasing the population of premature infants due to improving neonatal care we try to find other independent criteria in addition to birth weight and gestational age to reduce the number of infants who need a retinopathy of prematurity (ROP) screening examination.

Methods: This is a retrospective cross-sectional study, included 150 preterm infants with gestational age of ≤ 32 weeks or birth weight ≤ 1500 g or receives oxygen therapy for more than 2 days, who were examined from the fourth week of life for ROP in Tabriz Al-Zahra hospital's neonatal intensive care unit (NICU), from March 2017 to January 2018. All infants with other risk factors such as multi gestational pregnancy, blood transfusion or exchange, phototherapy, underlying disease, cerebral hemorrhage, chronic lung disease and sepsis were excluded from the study. Infants divided to two group ROP ($n=47$) and non-ROP ($n=133$) and their 5-minute Apgar score was compared as an independent risk factor. Exploratory data analysis was performed using descriptive measures with independent samples t tests. Kolmogorov-Smirnov tests were used to test the normality of data distribution.

Results: The mean 5-minute Apgar scores were statistically lower in ROP group (5.4 ± 2.3) than non-ROP group (6.2 ± 2.1) (P value=0.041).

Conclusion: Apgar score was statistically significant, but in light of the very close results obtained, it seems prudent to consider an independent risk factor.

Introduction

Retinopathy of prematurity (ROP) is a multifactorial disorder with vasoproliferative nature that affecting premature neonates with low birth weight (LBW) and very low birth weight (VLBW). ROP is considered the main cause of preventable but untreatable blindness in children throughout the world.¹⁻⁴

Many risk factors can lead to the development of ROP; their leaders are gestational age and birth weight. Other factors described in the articles are oxygen therapy, male gender, apnea, sepsis, cerebral hemorrhages and anemia. The role of these risk factors has not yet been fully elucidated. Instead, preeclampsia and respiratory system maturation due to prenatal steroids therapy have been shown to have a protective role.⁵⁻⁹

Since the survival rate of premature infants' increases, recent articles have shown a significant decrease in the incidence of ROP, which is possible to improved neonatal

care and better determining the causes of ROP.

Identifying other effective risk factors and prognostic factors which can be easily evaluated can reduce the number of unnecessary examinations.

Because we observed cases of ROP in routine examinations of neonates who had no risk factor for screening we concluded that other measurable risk factors other than gestational age and weight should be considered for screening. Perhaps the first measurable risk factor that predicts the probability of oxygen therapy is the Apgar score. Our first objective in this study was to determine whether the Apgar score is an independent risk factor or not and in the following of this study, what score can be considered as a risk factor.

Methods

This retrospective cross-sectional study was performed from March 2017 to January 2018. The study population

*Corresponding Author: Ali Kiavar, Department of Ophthalmology, Tabriz University of Medical Sciences, Tabriz, Iran.

, alikiavar@gmail.com

© 2020 The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

consisted of newborns admitted to the neonatal intensive care unit (NICU) at Al-Zahra hospital of Tabriz. For the cases of gestational age ≤ 32 weeks or birth weight ≤ 1500 g or receive oxygen therapy for more than 2 days, then screening for ROP started between 4 and 6 weeks after birth and re-examined from 7 to 14 days according to the fundus exam until complete vascularization of the retina was obtained also treatment was performed in all cases with type 1 ROP.

A single drop of tropicamide (1%) and phenylephrine (2.5 mg/mL) was instilled into each eye to obtain full mydriasis 1 hour and 30 minutes before starting examination. An ROP expert investigator examined the fundus using a Topcon indirect ophthalmoscope and a VOLK Pan Retinal 2.2D lens in NICU of Alzahra hospital or Nikukary eye center if permission from NICU. If retinal vascularization was completed without any stage of ROP, then the infant was registered to the non-ROP group. Upon the witness of any stage of ROP, the infant was registered to ROP group.

These parameters were examined and recorded: gestational age, gender, birth weight, duration and method of treatment with oxygen, Apgar score at 5 minute. All infants with the following characteristics were excluded from the study: multi gestational pregnancy, blood transfusion or exchange, phototherapy, underlying disease, cerebral hemorrhage, hypothermia, chronic lung disease, sepsis, jaundice, pulmonary hemorrhage and pneumothorax also infants who did not receive regular follow-up until complete retinal vascularization.

SPSS, version 23 was used for statistical analysis. Descriptive statistics were performed with independent samples t tests and Kolmogorov-Smirnov tests were used to assess the normality of data distribution. The coefficient level of this study was 0.95 and a difference with $P < 0.05$ was considered statistically significant.

Results

One hundred and sixty-five neonates have inclusion criteria but 15 infants excluded because their families change their location. Finally, 150 neonates were included in the study. 47 infants (31.3%) developed ROP in their examination and 103 infants (68.7%) never developed any stage of ROP. The mean gestational ages in the ROP and non-ROP groups were 27.83 ± 1.70 weeks and 29.07 ± 1.55 weeks, respectively (P value < 0.001). The mean birth weight in the ROP and non-ROP groups were 1180.53 ± 495.12 g and 1079.45 ± 190.45 grams, respectively (P value = 0.64) and the mean Apgar scores were 5.4 ± 2.3 in ROP group and 6.2 ± 2.1 in non-ROP group, respectively (P value = 0.041) (Table 1).

Discussion

Gestational age and birth weight are traditionally considered the main risk factors for development of ROP; they used as screening criteria all over the world.

Table 1. Neonatal Characteristics of ROP and Non-ROP groups.

	Non-ROP (n=103)	ROP (n=47)	P value
Age	29.54 \pm 5.73	30.79 \pm 5.32	0.209
Birth weight (g)	1079.45 \pm 190.45	1180.53 \pm 495.70	0.641
Gestational age (wk)	29.07 \pm 1.55	27.83 \pm 1.70	<0.001
Duration of oxygen therapy (day)	3.22 \pm 2.62	5.78 \pm 3.57	0.022
Apar score	6.19 \pm 2.04	5.43 \pm 2.28	0.041

recommend criteria for ROP screening were; infants with a birth weight of ≤ 1500 g or gestational age of ≤ 30 weeks and with a birth weight of 1500–2000 grams or gestational age of > 30 weeks with an unstable clinical course.¹⁰

It must be consider that these criteria vary in different countries depending on socioeconomic conditions, such as in our country (I.R. Iran) these limits are birth weight ≤ 2000 g and gestational age ≤ 34 weeks for screening of ROP.

Saving of more immature born neonates leads to increases the population of infants in need of ophthalmological screening, thus use other criteria can reduce inessential aggressive fundus examinations of low risk infants and consequently economical savings occur.

In a study of 957 infants in 2011, Ke et al also found the low Apgar score at 5 minutes was an effective risk factor for the ROP progression to stages requiring treatment. But they could not prove it as an independent factor.¹¹

Abrishami et al, in a study in Mashhad with 122 cases founded that Apgar score in ROP cases were lower than newborns without ROP but their study was multifactorial and they not isolated Apgar score as independent factor.¹²

Another study done with Marinov et al in Bulgaria, they shown A low 5-minute Apgar score and an Apgar score of six or less at 5 minutes were not statistically significant risk factors for increasing the incidence of ROP, but were significant risk factor for the progression of low stage ROP to stages requiring treatment.¹³

Yang et al who conducted a retrospective study of 252 (VLBW) children (birth weight < 1500 g) found that low 5-minute Apgar score was an important risk factor for development of ROP requiring treatment.¹⁴

Our study shown that 5-minute Apgar score was statistically an independent risk factor for the incidence of ROP but given the very statistically close results, it may not be very strong criteria for screening of ROP but it is a good predictor for progression to ROP.

Our study has shortcomings that need further studies to complete; we input any stage of ROP in case group but due to results it seems best to evaluate the different stage of ROP separately. Also in our study gestational age ≤ 32 weeks was a risk factor for ROP but in infants with birth weight ≤ 1500 g, birth weight not considered as a risk factor for ROP.

Conclusion

Today, due to the great advances in the field of NICU and neonatology care, more (VLBW) and extremely low birth weight infants survive so that the infants that need ROP screening are increasing and we should find additional criteria for decreasing the population of whom needs screening our study shown Apgar 5 score is risk factor for ROP but statically not a strong criteria for reducing screening population.

Conflict of Interest

There is no conflict of interest.

Ethic approval

This research was approved by regional ethic committee of Tabriz University of medical sciences(IR.TBZMED. REC.1397.959)

Acknowledgment

This study was financially supported by Tabriz University of Medical Sciences.

Authors' Contribution

A.E.M. carried out the design and coordinated the study, participated in fundus exams. M.R.N. participated in neonate's examinations and follow ups. A.K. provide assistance in the design of the study, neonate's examinations and follow ups. H.S. provided assistance in statistical analysis and manuscript preparation. A.M.F and Z.A. assisted in data gathering and participated in manuscript editing.

References

1. Quinn GE, Gilbert C, Darlow BA, Zin A. Retinopathy of prematurity: an epidemic in the making. *Chin Med J (Engl)*. 2010;123(20):2929-37.
2. Gilbert C. Retinopathy of prematurity: a global perspective of the epidemics, population of babies at risk and implications for control. *Early Hum Dev*. 2008;84(2):77-82. doi: 10.1016/j.earlhumdev.2007.11.009.
3. Gilbert C, Fielder A, Gordillo L, Quinn G, Semiglia R, Visintin P, et al. Characteristics of infants with severe retinopathy of prematurity in countries with low, moderate, and high levels of development: implications for screening programs. *Pediatrics*. 2005;115(5):e518-25. doi: 10.1542/peds.2004-1180.
4. Abdel HA, Mohamed GB, Othman MF. Retinopathy of

- prematernity: a study of incidence and risk factors in NICU of Al-Minya University Hospital in Egypt. *J Clin Neonatol*. 2012;1(2):76-81. doi: 10.4103/2249-4847.96755.
5. Horbar JD, Carpenter JH, Badger GJ, Kenny MJ, Soll RF, Morrow KA, et al. Mortality and neonatal morbidity among infants 501 to 1500 grams from 2000 to 2009. *Pediatrics*. 2012;129(6):1019-26. doi: 10.1542/peds.2011-3028.
6. Coats DK. Retinopathy of prematurity: Pathogenesis, epidemiology, classification, and screening. Available from: <http://www.uptodate.com/contents/retinopathy-of-prematurity>. Updated July 27, 2015. Accessed July 2015.
7. Hellström A, Smith LE, Dammann O. Retinopathy of prematurity. *Lancet*. 2013;382(9902):1445-57. doi: 10.1016/s0140-6736(13)60178-6.
8. Chawla D, Agarwal R, Deorari A, Paul VK, Chandra P, Azad RV. Retinopathy of prematurity. *Indian J Pediatr*. 2012;79(4):501-9. doi: 10.1007/s12098-010-0279-7.
9. Filho JB, Eckert GU, Valiatti FB, da Costa MC, Bonomo PP, Procianny RS. Prevalence and risk factors for retinopathy of prematurity: study with 450 very low birth weight preterm infants. *Rev Bras Oftalmol*. 2009;68(1):22-9. doi: 10.1590/S0034-72802009000100005.
10. Section on Ophthalmology, American Academy of Pediatrics, American Academy of Ophthalmology, American Association for Pediatric Ophthalmology and Strabismus. *Pediatrics*. 2006;117(2):572-6. doi: 10.1542/peds.2006-0343.
11. Ke XY, Ju RH, Zhang JQ, Chen H, Wei EX, Chen XH. Risk factors for severe retinopathy of prematurity in premature infants: a single-center study. *Nan Fang Yi Ke Da Xue Xue Bao*. 2011;31(12):1963-7.
12. Abrishami M, Maemori GA, Boskabadi H, Yaeghobi Z, Mafi-Nejad S, Abrishami M. Incidence and risk factors of retinopathy of prematurity in Mashhad, northeast Iran. *Iran Red Crescent Med J*. 2013;15(3):229-33. doi: 10.5812/ircmj.4513.
13. Marinov VG, Koleva-Georgieva DN, Sivkova NP, Krasteva MB. The 5-minute Apgar score as a prognostic factor for development and progression of retinopathy of prematurity. *Folia Med (Plovdiv)*. 2017;59(1):78-83. doi: 10.1515/folmed-2017-0011.
14. Yang CY, Lien R, Yang PH, Chu SM, Hsu JF, Fu RH, et al. Analysis of incidence and risk factors of retinopathy of prematurity among very-low-birth-weight infants in North Taiwan. *Pediatr Neonatol*. 2011;52(6):321-6. doi: 10.1016/j.pedneo.2011.08.004.